Central Illinois Natural Health Clinic 1012 W. Fairchild Street • Danville IL, 61832 (217) 443-4372

Oswestry Low Back Pain Disability Questionnaire

Name __	Date
	Please mark the ONE choice from EACH group that best describes you.
PAIN	INTENSITY
$\Box A$.	The pain comes and goes and is very mild.
□ B.	The pain is mild and does not vary much.
\Box C.	The pain comes and goes and is moderate.
□D.	The pain is moderate and does not vary much.
□E.	The pain comes and goes and is severe.
□F.	The pain is severe and does not vary much.
PERS	ONAL CARE
$\Box A$.	I would not have to change my way of washing or dressing in order to avoid pain.
□ B.	I do not normally change my way of washing or dressing even though it causes some pain.
\Box C.	Washing and dressing increases the pain, but I manage not to change my way of doing it.
□D.	Washing and dressing increases the pain and I find it necessary to change my way of doing it.
□E.	Because of the pain, I am unable to do some washing and dressing without help.
□F.	Because of the pain, I am unable to do any washing or dressing without help.
LIFTING	
$\Box A$.	I can lift heavy weights without extra pain.
□ B.	I can lift heavy weights, but it causes extra pain.
\Box C.	Pain prevents me from lifting heavy weights off the floor.
□D.	Pain prevents me from lifting heavy weights off the floor, but I can manage if they are
	conveniently positioned, e.g., on a table.
□E.	Pain prevents me from lifting heavy weights, but I can manage light to medium weights if
	they are conveniently positioned.
□ F.	I can only lift very light weights, at the most.
WALKING	
$\Box A$.	Pain does not prevent me from walking any distance.
□ B.	Pain prevents me from walking more than one mile.
\Box C.	Pain prevents me from walking more than ½ mile.
\Box D.	Pain prevents me from walking more than 1/4 mile.
□E.	I can only walk while using a cane or on crutches.
□F.	I am in bed most of the time and have to crawl to the toilet.
SITTI	
\Box A.	I can sit in any chair as long as I like without pain.
□ B.	I can only sit in my favorite chair as long as I like.
\Box C.	Pain prevents me from sitting more than one hour.
□D.	Pain prevents me from sitting more than ½ hour.
□E.	Pain prevents me from sitting more than ten minutes.
□F.	Pain prevents me from sitting at all.

Patient SignatureDate	
CHAN A. B. C. D. E.	My pain is rapidly getting better. My pain fluctuates, but overall is definitely getting better. My pain seems to be getting better, but improvement is slow at present. My pain is neither getting better nor worse. My pain is gradually worsening. My pain is rapidly worsening.
□D. □E. □F.	I get extra pain while traveling which compels me to seek alternative forms of travel. Pain restricts all forms of travel. Pain prevents all forms of travel except that done lying down.
TRAV □A. □B. □C.	I get no pain while traveling. I get some pain while traveling, but none of my usual forms of travel make it any worse. I get extra pain while traveling, but it does not compel me to seek alternative forms of travel.
□D. □E. □F.	interests, e.g., dancing, etc. Pain has restricted my social life and I do not go out very often. Pain has restricted my social life to my home. I have hardly any social life because of the pain.
SOCL □ A. □ B. □ C.	AL LIFE My social life is normal and gives me no pain. My social life is normal, but increases the degree of my pain. Pain has no significant effect on my social life apart from limiting my more energetic
SLEE □ A. □ B. □ C. □ D. □ E. □ F.	I get no pain in bed. I get pain in bed, but it does not prevent me from sleeping well. Because of pain, my normal night's sleep is reduced by less than one-quarter. Because of pain, my normal night's sleep is reduced by less than one-half. Because of pain, my normal night's sleep is reduced by less than three-quarters. Pain prevents me from sleeping at all.
□A. □B. □C. □D. □E. □F.	I can stand as long as I want without pain. I have some pain while standing, but it does not increase with time. I cannot stand for longer than one hour without increasing pain. I cannot stand for longer than ½ hour without increasing pain. I cannot stand for longer than ten minutes without increasing pain. I avoid standing, because it increases the pain straight away.

STANDING